

AD-A062 005

MASSACHUSETTS INST OF TECH CAMBRIDGE DEPT OF CHEMISTRY F/G 20/5  
FORMATION AND DEACTIVATION PROCESSES IN ELECTRONIC TRANSITIONS,--ETC(U)  
OCT 78 J I STEINFELD AFOSR-75-2758

UNCLASSIFIED

AFOSR-TR-78-1509

NL

1 OF 1  
AD  
A062005



END  
DATE  
FILMED  
3-79  
DDC

ADA062005

DDC FILE COPY

AFOSR-TR- 78-1509

FINAL REPORT

to the

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH

on

Grant No. AFOSR-75-2758

"Formation and Deactivation Processes in  
Electronic Transitions, Chemically Pumped Lasers"

by

Jeffrey I. Steinfeld  
Associate Professor of Chemistry  
Massachusetts Institute of Technology  
Cambridge, Massachusetts 02139

(1 October 1978)

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)  
NOTICE OF TRANSMITTAL TO DDC  
This technical report has been reviewed and is  
approved for public release (AR 190-12 (7b)).  
Distribution is unlimited.  
A. D. BLOSE  
Technical Information Officer

Approved for public release;  
distribution unlimited.

78 12 04.065



The present research program was concerned with measurement of the rates of formation of excess populations of electronically excited small molecules, particularly in chemiluminescent metal + oxidant reactions, and characterization of the deactivation processes which tend to relax these excited states. One of the motivations for these investigations was the possibility of finding a suitable candidate for a chemically - pumped, electronic transition (i.e., visible - wavelength) laser system. Although such a system was not developed in the course of this work, a great deal of information on the pertinent reaction and relaxation rates was obtained. Initially, this program was under the joint direction of Profs. J.L. Gole and J.I. Steinfeld at M.I.T.; Prof. Gole moved his part of the research to the Georgia Institute of Technology during the final year of the grant. For convenience, we summarize here the principal findings arising from the overall research program; more details can be found in the cited reference(s) from the publications list which follows.

1. Aluminum - Ozone Reaction. The reaction of aluminum with ozone produces the  $B^2\Sigma^+ \rightarrow X^2\Sigma^+$  emission spectrum of AlO. Coupling between the  $B^2\Sigma^+$  and  $A^2\Pi$  levels of AlO is attributed to perturbations (Lindsay and Gole, 1977).

2. Lanthanum - Oxidant Reactions. Reaction of lanthanum with oxidants ( $O_2$ ,  $NO_2$ ,  $N_2O$ ,  $O_3$ ) results in chemiluminescent emission from the  $A^2\Pi$ ,  $B^2\Sigma^+$ ,  $C^2\Pi$ , and in the latter three instances the  $D^2\Sigma^+$  states of LaO. The dissociation energy of LaO and a potential function for its  $D^2\Sigma^+$  state have been deduced from the data. The electronic state distribution

78 12 04. 065

in the  $\text{La} + \text{O}_2$  reaction has been shown by Levine and Ben-Shaul to follow a simple zero-surprisal form (Gole and Chalek, 1976).

3. Reactions of Silanes and Boranes with Oxidants. The reactive systems  $\text{B}_2\text{H}_6 + \text{O}_3, \text{F}_2$ ;  $\text{B}_5\text{H}_9 + \text{O}_3, \text{F}_2$ ; and  $\text{Si}_2\text{H}_6 + \text{O}_3, \text{F}_2$  have been studied. The  $\text{B}_2\text{H}_6 + \text{O}_3$  and  $\text{B}_5\text{H}_9 + \text{O}_3$  systems yield primarily  $\text{BO}_2^*$  with some  $\text{BO}^*$ . The  $\text{B}_2\text{H}_6 + \text{F}_2$  and  $\text{B}_5\text{H}_9 + \text{F}_2$  systems yield  $\text{HBF}^*$  and  $\text{BH}^*$ . The  $\text{Si}_2\text{H}_6 + \text{O}_3$  system is believed to yield  $\text{HSiO}$ . The reaction of disilane ( $\text{Si}_2\text{H}_6$ ) with  $\text{F}_2$  is characterized by strong emission from the  $\text{A}^2\Sigma^+ - \text{X}^2\Pi$  transition of  $\text{SiF}$ , the  $\text{A}(^2\text{B}_1) - \text{X}(^2\text{A}_1)$  transition of  $\text{SiF}_3$ , and the  $\text{A}^2\Delta - \text{X}^2\Pi$  transition of  $\text{SiH}$ . By contrast, the principal emission from monosilane - chlorine or disilane - chlorine reactions is the  $^1\text{A}'' - ^1\text{A}'$  transition of  $\text{HSiCl}$ . An attempt was made to run the  $\text{B}_5\text{H}_9 + \text{F}_2$  reaction in a laser cavity, but stimulated emission was not observed (Conner et al., 1977).

4. Scandium and Yttrium Oxidation Reactions. The reactions of Sc and Y with  $\text{O}_2$  are characterized by emission from the  $\text{A}^2\Pi$  and the previously unobserved  $\text{A}^1^2\Delta$  states of  $\text{ScO}$  and  $\text{YO}$ . The reactions of Sc and Y with  $\text{NO}_2, \text{N}_2\text{O}$  and  $\text{O}_3$  result in emission from the  $\text{A}^2\Pi$  and  $\text{B}^2\Sigma^+$  states of  $\text{ScO}$  and  $\text{YO}$ . Dissociation energies of  $\text{ScO}$  and  $\text{YO}$  [ $\text{D}_0^0(\text{YO}) \geq 7.49 \text{ eV}$ ] have been deduced from the data (Chalek and Gole, 1977).

5. Titanium-Oxidant Reactions. The reaction of Ti with  $\text{O}_2, \text{NO}_2$ , and  $\text{N}_2\text{O}$  results in chemiluminescent emission from the  $\text{B}^3\Pi$  and  $\text{C}^3\Delta$  states. From analysis of the temperature dependence, we conclude that  $\text{Ti}(^5\text{F})$  excited atoms are the most likely precursors of the  $\text{C}^3\Delta$  state (Dubois and Gole, 1977).

ACCESSION for	
NTIS	White Section <input checked="" type="checkbox"/>
DDC	B.N. Section <input type="checkbox"/>
UNANNOUNCED	
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
SPECIAL	
A	



6. Chemiluminescent Reactions of the Group IIIB Metals. The reactions of scandium, yttrium, and lanthanum with  $F_2$ ,  $Cl_2$ ,  $Br_2$ ,  $SF_6$ ,  $ClF$ , and  $IBr$  have been studied under "single collision" conditions using a beam-gas arrangement. The observed chemiluminescent spectra are primarily attributed to emission from the metal monohalides. Dihalide emission is also observed in the Sc-Br and Sc-Cl systems. We find extremely selective production of monohalide excited electronic states in both the scandium and yttrium reactions. The lanthanum reactions are characterized by relatively moderate selective excitation of monohalide excited states. The selective feature dominating the scandium and yttrium monohalide spectra has not been observed by previous researchers. On the basis of molecular orbital theory, energy conservation, and known spectra for scandium and yttrium fluoride, the selective emission feature is assigned to a  $^3\Sigma^+ - ^1\Sigma^+$  or  $^3\Sigma^+ - ^3\Delta$  band system. We have determined the molecularity with respect to oxidant for all reactions studied, monitoring chemiluminescent intensity versus oxidant pressure. All reactions producing metal monofluoride are found to be first order with respect to oxidant while reactions producing the monochloride or monobromide are found to be "faster than first order." This behavior is explained in terms of the kinetics of metastable excited states. The production of  $ScCl_2$  and  $ScBr_2$  also occurs via a "faster than first order" process. Based upon mechanistic considerations entailed in the kinetics of metastable excited states, the radiative lifetimes of the selectively populated excited states of  $ScCl$ ,  $ScBr$ , and  $YCl$  are estimated to be on the order of  $2 \times 10^{-3}$  seconds. Analysis of the temperature dependence for six representative reactions indicates that selective

excited state formation of the metal monohalides proceeds by a direct mechanism with negligible activation energy. From the short wavelength limits of the chemiluminescent spectra, lower bounds to the dissociation energies of the monohalides have been determined. For some of the chemiluminescent reactions studied, we have found evidence for "super-relaxation" on collision.

In the reactions of Group IIIB metals with  $F_2$  and  $NO_2$ , cross sections for excited-state metal monohalide formation considerably exceed those for monoxide formation. The experimentally determined cross sections are compared with total reactive cross sections, calculated on the basis of mechanistic models used to describe the oxidation reactions of Group IIIB atoms. Based upon this comparison, the ratio of excited state to ground state formation (quantum yield) is determined to be not less than 0.1 for the monohalides (Chalek *et al.*, 1976; Gole *et al.*, 1977; Preuss and Gole, 1977).


7. Optical Model for Surface Crossing. Several approaches were investigated to the problem of calculating the rates of collision-induced relaxation between molecular electronic states. Following a classical trajectory ("surface-hopping") study of  $I_2^* B^3\Pi_{ou} \rightarrow ^3\Pi_g$  (dissociative), the same system was analyzed using an equivalent optical-potential model which yielded a great saving in computation time. This model was also applied to the  $N_2^* B^3\Pi \rightarrow (A^3\Sigma, W^3\Delta)$  system, but difficulties were encountered with  $\Omega \neq 0$  systems, such as  $CN A^2\Pi$ . The Landau-Zener calculation of the probabilities employed in the optical model was also compared with more recent semiclassical scattering theories (Garetz *et al.*, 1975, 1976; Bendazzoli *et al.*, 1977).

8. Information-Theoretic Analysis of Deactivation Rates. The Information-Theoretic treatment of nonequilibrium processes, due to Levine and Bernstein, has proven to be remarkably useful for characterizing relaxation processes, and several aspects of applying these methods were investigated in this program. A set of state-to-state rates for V-V relaxation in HF was calculated by the "Surprisal Synthesis" method (work performed at Air Force Weapons Laboratory, Kirtland AFB), and the nonlinear master equation governing V-V equilibration in HF was solved using exponential-gap rate constants, in collaboration with Professor R.D. Levine (Steinfeld, 1978; Tabor et al., in press).

9. Spectroscopy and Deactivation of CN  $A^2\Pi$ . Laser action has been observed from the CN A state produced by photodissociation or electric-discharge pumping, and this system has also been suggested as a promising candidate for a chemically-pumped laser system. Since little is known about the quenching of this state, several investigations were undertaken. Some 39 bands of the "Berkeley" atlas of the CN  $A^2\Pi - X^2\Sigma$  transition, along with 16 bands measured at high resolution by R. Bacis of the University of Lyon, were reanalyzed with a least-squares technique, allowing for all ( $A^2\Pi \sim X^2\Sigma$ ) and ( $A^2\Pi \sim B^2\Sigma$ ) perturbations. This provided mixing coefficients for the perturbed levels in the A state. The analysis is currently being completed in collaboration with Dr. J.A. Coxon, Dalhousie University, Halifax, Nova Scotia. Attempts to carry out monochromatically excited fluorescence experiments on selected levels of the CN A state were, however, inconclusive (Kotlar et al., to be published).



10. Near-Resonant Electronic Energy Transfer. In collaboration with Dr. D. Sutton of the Aerospace Corporation, a measurement was carried out of the rate constant for the nearly resonant ( $\Delta E = 16 \text{ cm}^{-1}$ ) energy transfer process,



The measured value,  $1 \times 10^{-9} \text{ cm}^3 \text{ molecule}^{-1} \text{ sec}^{-1}$  ( $\sigma \approx 200 \text{ \AA}$ ), is one of the largest that has been observed for such a process. Since both optical transitions are highly forbidden, a simple dipole interaction model cannot account for this observation. A technically important example of such a process is the chemically pumped  $\text{O}_2^*(^1\Delta) - \text{I}$  atom transfer laser recently developed at AFWL (Capelle et al., in press).

11. Multiphoton  $\text{CO}_2$  Laser Vibrational Heating of Metastable Electronic States. Biacetyl molecules, prepared in the triplet metastable state by intersystem crossing from the  $^1\text{A}_g$  electronic state which in turn is excited by irradiation with the 457.9 nm line of a cw  $\text{Ar}^+$  laser, are pumped by an intense  $\text{CO}_2$  laser pulse which excites the system vibrationally. Changes in the luminescence characteristics of the electronically excited molecules induced by the infrared pulse are studied in various visible spectral regions. In the region characteristic of the biacetyl phosphorescence, a fast decay of the phosphorescence is observed. A burst of delayed fluorescence is observed in the fluorescence region of the biacetyl; this signal is characterized by a very fast rise time. Both fluorescence and phosphorescence signal exhibit the same decay time.



Attenuation of the CO<sub>2</sub> laser beam results in an increase of the decay time and in a decrease of the amplitudes of the fluorescence signal.

The experimental results are explained in terms of mixed singlet-triplet vibronic wave functions, characterizing the triplet vibrational manifold, with vibrational energy exceeding the  $^1A_u - ^3A_u$  energy separation. A calculation of the decay times as a function of the absorbed CO<sub>2</sub> laser energy, which is based on a recent model for delayed fluorescence, gives good agreement with the experimental data. This is the first well-characterized example of reverse intersystem crossing resulting from multiple infrared photon excitation of vibrational modes, which are then converted into electronic excitation (Burak *et al.*, in press; preprint enclosed with this report).

### List of Publications

B. Garetz, L.L. Poulsen, and J.I. Steinfeld, "Optical Model for Potential Surface Crossing," *Chem. Phys.* 9, 385 (1975).

B. Garetz, J.I. Steinfeld, and L.L. Poulsen, "Collision-Induced Electronic Deactivation of  $N_2^*(B^3\Pi_g)$ ," *Chem. Phys. Letters* 38, 365 (1976).

C. Chalek, D.R. Preuss, and J.L. Gole, "Single Collision Selective Chemical Reaction of Group III Metals (Sc, Y, La) and Halogens ( $F_2$ ,  $ClF$ ,  $SF_6$ , and  $Cl$ )," in Proceedings, 2nd Summer Colloquium on Electronic Transition Lasers, (J.I. Steinfeld, ed.), p. 50, M.I.T. Press, Cambridge, Mass., 1976.

J.L. Gole, D.R. Preuss, and C.L. Chalek, "Kinetics of Metastable Excited State Products in a Beam-Gas Chemiluminescent Reaction," *J. Chem. Phys.*, 66, 548 (1977).

D.M. Lindsay and J.L. Gole, " $Al + O_3$  Chemiluminescence: Perturbations and Vibrational Population Anomalies in the  $B^2\Sigma^+$  State of  $AlO$ ," *J. Chem. Phys.*, 66, 3886 (1977).

J.L. Gole and C.L. Chalek, "Characterization of the Ground and Excited States of Lanthanum Oxide through Bimolecular Oxidation of La Metal with  $O_2$ ,  $NO_2$ ,  $N_2O$ , and  $O_3$ ," *J. Chem. Phys.*, 65, 4384 (1976).

C.P. Conner, G.W. Stewart, D.M. Lindsay, and J.L. Gole, "Reactions of Silanes with Halogens: Chemiluminescent Products in the UV-Visible Spectrum," *J. Am. Chem. Soc.* 99, 2540 (1977).

C.L. Chalek and J.L. Gole, "Single Collision Chemiluminescence Studies of Scandium and Yttrium Oxidation with  $O_2$ ,  $NO_2$ ,  $N_2O$ , and  $O_3$ ," *Chem. Phys.*, 19, 59 (1977).

L.H. Dubois and J.L. Gole, "Bimolecular Single Collision Reaction of Ground and Metastable Excited States of Titanium with  $O_2$ ,  $NO_2$ , and  $N_2O$  - Confirmation of  $D_0(TiO)$ ," *J. Chem. Phys.*, 66, 779 (1977).

J.L. Gole, "Development of Visible Chemical Lasers from Reactions Yielding Visible Chemiluminescence," in Proceedings, 3rd Summer Colloquium on Electronic Transition Lasers (J.I. Steinfeld, S. Suchard, and L.E. Wilson, eds.) p. 136, M.I.T. Press, Cambridge, Mass., 1977.

D.R. Preuss and J.L. Gole, "The Temperature Dependence of Chemiluminescent Reactions - Activation Energies for Excited State Formation," in Proceedings, 3rd Summer Colloquium on Electronic Transition Lasers (J.I. Steinfeld, S. Suchard, and L.E. Wilson, eds.), p. 176, MIT Press, Cambridge, Mass., 1977.

J.L. Gole, "High Temperature Chemistry: Modern Research and New Frontiers," *Ann. Rev. Phys. Chem.* 27, 525 (1976).

J.I. Steinfeld, "Fluorescence and Double Resonance Studies of Energy Transfer," in Proceedings, Conference on Energy Transfer Processes in Chemical Reactions, Konigstein, Sept. 1976; published in Berichte der Deutsche Bunsengesellschaft 81, 221 (1977).

G.L. Bendazzoli, M. Ruimondi, B.A. Garetz, T.F. George, and K. Morokuma, "Semiclassical Study of Collision Induced Predissociation: Comparison of the Landau-Zener Model with the Method of Analytic Continuation", Theoretica Chimica Acta, 44, 341 (1977).

J.L. Gole, D.R. Preuss, and C.L. Chalek, "Kinetics of Metastable Excited State Products in a Beam-Gas Chemiluminescent Reaction," J. Chem. Phys. 66, 548 (1977).

J.I. Steinfeld, "Activation and Deactivation Rates in High-Power Lasers", Proceedings, Colloquium on High-Power Lasers and Applications (Munich, June 1976); Series in Optical Sciences No. 9 (K.-L. Kompa, and H. Walther, eds.), pp. 66-73, Springer-Verlag, Berlin, 1978.

D.R. Preuss and J.L. Gole, "The Use of Temperature Dependent Reaction Rates to Correct Metal Halide (Oxide) Dissociation Energies as Determined from Chemiluminescence Reactions", J. Chem. Phys. 66, 880 (1977).

M. Tabor, R.D. Levine, A. Ben-Shaul, and J.I. Steinfeld, "Microscopic and Macroscopic Analysis of Non-Linear Master Equations: Vibrational Relaxation of Polyatomic Molecules", Mol. Phys., in press.

G.A. Capelle, D.G. Sutton, and J.I. Steinfeld, "Near-Resonant Electronic Energy Transfer from  $\text{NF}(a^1\Delta)$  to Bi", J. Chem. Phys., in press.

I. Burak, T.J. Quelly, and J.I. Steinfeld, "Multiphoton  $\text{CO}_2$  Laser Vibrational Heating of the Triplet State of Biacetyl", J. Chem. Phys., in press.

A.J. Kotlar, R.W. Field, and J.I. Steinfeld, "Analysis of Perturbations in the  $\text{A}^2\Pi - \text{X}^2\Sigma^+$  'Red' System of CN", to be published.

J.I. Steinfeld, "Formation and Deactivation Processes in Electronic Transitions, Chemically Pumped Lasers", Final Technical Report.



REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER <b>AFOSR-TR-78-1509</b>	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER <i>Final rept. 1 Sep 74 - 30 Sep 78</i>
4. TITLE (and Subtitle) <b>Formation and Deactivation Processes in Electronic Transition; Chemistry Pumped Lasers</b>	5. TYPE OF REPORT & PERIOD COVERED <b>Final; 9/1/74-9/30/78</b>	
6. AUTHOR(s) <b>Jeffrey I. Steinfeld</b>	7. PERFORMING ORG. REPORT NUMBER <b>15</b>	8. CONTRACT OR GRANT NUMBER(s) <b>AFS0R-75-2758</b>
9. PERFORMING ORGANIZATION NAME AND ADDRESS <b>Department of Chemistry Massachusetts Institute of Technology Cambridge, Massachusetts 02139</b>	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS <b>61102F 2303/B1</b>	
11. CONTROLLING OFFICE NAME AND ADDRESS <b>Air Force Office of Scientific Research /NC Bolling AFB Washington, D.C. 20332</b>	12. REPORT DATE <b>10/1/78</b>	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) <b>11 1 Oct 78 12 12p.</b>	13. NUMBER OF PAGES <b>10</b>	
	15. SECURITY CLASS. (of this report) <b>UNCLASSIFIED</b>	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report)  <b>Approved for public release; distribution unlimited</b>		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) <b>Chemical Lasers      Energy Transfer Chemiluminescence      Quantum Yields Deactivation Rates      Relaxation</b>		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  <b>Research was performed on the rates of formation of excess populations of electronically excited small molecules, particularly in chemiluminescent metal + oxidant reactions, and characterization of the deactivation processes which tend to relax these excited states. Results were obtained in the following areas: (1) aluminum + ozone reaction; (2) lanthanum + oxidant reactions; (3) reactions of silanes and boranes with oxidants; (4) scandium and yttrium oxidant reactions; (5) titanium + oxidant reaction; (6)</b>		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

220 007  
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

chemiluminescent reactions of the Group IIB metals; (7) optical model for surface crossing; (8) Information-Theoretic analysis of deactivation rates; (9) Spectroscopy and deactivation of  $\text{CN}(A^2\Pi)$ ; (10) near-resonant electronic energy transfer; and (11) multiphoton  $\text{CO}_2$  laser vibrational heating of metastable electronic states.

$A_1 2 \Pi_i$

UNCLASSIFIED